



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Now $a < h$, $b < k$. $\therefore ah + bk < h^2 + k^2$.

Calling $ah + bk$, r , we have $r^2 + 1$, and therefore $r^2 + 1 + s(r + s)$, or $r^2 + rs + s^2 + 1$, divisible by $r + s$.

It is evident that the number $ah + bk$ which is less than $h^2 + k^2$ and is such that $(ah + bk)^2 + 1$ is divisible by $h^2 + k^2$, can always be found.

Also solved by G. B. M. Zerr.

PROBLEMS FOR SOLUTION.

ALGEBRA.

303. Proposed by PROF. R. D. CARMICHAEL, Anniston, Ala.

Evaluate the determinant

$$\begin{vmatrix} D_1 & x_1x_2 & x_1x_3 & \dots & x_1x_n \\ x_1x_2 & D_2 & x_2x_3 & \dots & x_2x_n \\ x_1x_3 & x_2x_3 & D_3 & \dots & x_3x_n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ x_1x_n & x_2x_n & x_3x_n & \dots & D_n \end{vmatrix}$$

304. Proposed by C. N. SCHMALL, New York City.

A policeman on a motor-cycle starts in pursuit of an automobile when the latter has a headway of $\frac{1}{2}$ a mile. A pedestrian who is $\frac{1}{4}$ of a mile ahead of the auto and who is walking at the rate of 5 miles an hour, notices that when the auto overtakes him the policeman is only $\frac{5}{12}$ of a mile behind the auto, and $2\frac{1}{2}$ miles from where the officer started; he overtakes the auto. How long did the chase last?

305. Proposed by S. A. COREY, Hiteman, Iowa.

Prove or disprove, that $\sum_{n=1}^{n=\infty} \frac{1}{(2n-1)^2 + 4} = \frac{\pi}{4} - \frac{\pi}{8} \left(\frac{\cosh \pi}{\sinh \pi} \right)$.

GEOMETRY.

336. Proposed by F. H. HODGE, The University of Chicago.

A man owning a rectangular field $b=300$ feet by $a=600$ feet, wishes to lay out driveways of equal width having the diagonals of the field as center lines and such that the area of the driveways shall be $n/m=$ one-half, of the area of the field. Determine the width of the driveways.

337. Proposed by T. N. HILDEBRANT, The University of Chicago.

Required the locus of the vertices of the parabolae having a given focus and passing through a given point.

338. Proposed by C. N. SCHMALL, 239 East 7th Street, New York.

Given the base and vertical angle of a triangle, find the locus of the center of its "nine-point" circle. [Ex. 28, p. 65, Casey's *Sequel to Euclid*.]